

Explicit Toric Metric on Resolved Calabi-Yau Cone

Takeshi Oota^{1*} and Yukinori Yasui^{2†}

¹*Osaka City University, Advanced Mathematical Institute (OCAMI)*

3-3-138 Sugimoto, Sumiyoshi, Osaka 558-8585, Japan

²*Department of Mathematics and Physics, Graduate School of Science,*

Osaka City University

3-3-138 Sugimoto, Sumiyoshi, Osaka 558-8585, Japan

Abstract

We present an explicit non-singular complete toric Calabi-Yau metric using the local solution recently found by Chen, Lü and Pope. This metric gives a new supergravity solution representing D3-branes.

*e-mail: toota@sci.osaka-cu.ac.jp

†e-mail: yasui@sci.osaka-cu.ac.jp

D3-branes on the tip of toric Calabi-Yau cones have been extensively studied in connection with the AdS/CFT correspondence [1]. It is natural to consider the deformations of cone metrics in order to explore non-conformal theories [2, 3, 4, 5, 6, 7].

In this letter, we study a Calabi-Yau metric, i.e. Ricci-flat Kähler metric, constructed as the BPS limit of the six dimensional Euclideanised Kerr-NUT-AdS black hole metric [8]. In the black hole with equal angular momenta, the corresponding Calabi-Yau metric is of the form

$$g = \frac{(1-x)(1-z)}{3}(d\theta^2 + \sin^2 \theta d\phi^2) + \frac{(1-x)(x-z)}{f(x)}dx^2 + \frac{(1-z)(z-x)}{h(z)}dz^2 \quad (1)$$

$$+ \frac{f(x)}{9(1-x)(x-z)}(d\psi - \cos \theta d\phi + z(d\beta + \cos \theta d\phi))^2$$

$$+ \frac{h(z)}{9(1-z)(z-x)}(d\psi - \cos \theta d\phi + x(d\beta + \cos \theta d\phi))^2,$$

where

$$f(x) = 2x^3 - 3x^2 + a, \quad h(z) = 2z^3 - 3z^2 + b \quad (2)$$

with two parameters a and b . We assume that the roots x_i ($i = 1, 2, 3$) of $f(x) = 0$ are all distinct and real, and further they are ordered as $z_1 < x_1 < x_2 < x_3$ for the smallest real root z_1 of $h(z) = 0$. In order to avoid a curvature singularity we take the coordinates x and z to lie in the region $z \leq z_1 < x_1 \leq x \leq x_2 < 1$. Indeed, it is easy to see that such a singularity appears at $z = x$.

For $r \rightarrow \infty$ ($z = -r^2/2$), the metric tends to a cone metric $dr^2 + r^2 \bar{g}$, where

$$\bar{g} = \frac{1-x}{6}(d\theta^2 + \sin^2 \theta d\phi^2) + \frac{1-x}{2f(x)}dx^2 + \frac{f(x)}{18(1-x)}(d\beta + \cos \theta d\phi)^2 \quad (3)$$

$$+ \frac{1}{9}(d\psi - \cos \theta d\phi + x(d\beta + \cos \theta d\phi))^2.$$

This metric yields the Sasaki-Einstein metric $Y^{p,q}$ when we impose a suitable condition for the parameter a [9].

Next let us look at the geometry near $z = z_1$. We introduce new coordinates given by

$$\varphi_1 = -\frac{1}{2}(\psi - z_1\beta), \quad \varphi_2 = \frac{1}{1-z_1}(\psi + z_1\beta), \quad \varphi_3 = \phi. \quad (4)$$

Then the metric behaves as

$$g \simeq du^2 + u^2 d\varphi_1^2 + g^{(4)}, \quad (5)$$

where $u^2 = 2(z_1 - x)(z_1 - z)/(3z_1)$. Therefore, the periodicity of φ_1 should be 2π in order to avoid an orbifold singularity. The four dimensional Kähler metric $g^{(4)}$ is given by

$$g^{(4)} = \frac{(1-x)(1-z_1)}{3}(d\theta^2 + \sin^2 \theta d\varphi_3^2) + \frac{(1-x)(x-z_1)}{f(x)}dx^2 \\ + \frac{(1-z_1)^2 f(x)}{9(1-x)(x-z_1)}(d\varphi_2 - \cos \theta d\varphi_3)^2. \quad (6)$$

We now argue that by taking the special parameters $a = (1/2) - (1/32)\sqrt{13}$ which corresponds to $Y^{2,1}$, and $b = (1/4)(137 + 37\sqrt{13})$, the four dimensional space with metric $g^{(4)}$ is a non-trivial S^2 -bundle over S^2 , i.e. the first del Pezzo surface dP_1 . To see this, introduce a radial coordinate $y^2 = 2(x_i - z_1) |x - x_i| / |3x_i|$ on the (x, φ_2) -fibre space defined by fixing the S^2 coordinates θ and φ_3 in (6). Then, the fibre metric near boundary $x = x_i$ ($i = 1, 2$) is written as

$$dy^2 + \left(\frac{x_i(1-z_1)}{x_i - z_1} \right)^2 y^2 d\varphi_2^2. \quad (7)$$

Using the values of a and b , we have

$$x_1 = \frac{1}{8}(1 - \sqrt{13}), \quad x_2 = \frac{1}{8}(7 - \sqrt{13}), \quad z_1 = -\frac{1}{2}(2 + \sqrt{13}), \quad (8)$$

and then $x_i(1-z_1)/(x_i - z_1) = \pm 1/2$. The apparent singularities at $x = x_i$ can be avoided by choosing the periodicity of φ_2 to be 4π . Thus, the (x, φ_2) -fibre space is topologically S^2 . On the other hand, fixing the coordinate x in (6), we obtain a principal $U(1)$ -bundle over S^2 with the Chern number

$$\frac{1}{4\pi} \int_{S^2} d(-\cos \theta d\varphi_3) = 1. \quad (9)$$

The metric $g^{(4)}$ can be regarded as a metric on the associated S^2 -bundle of the principal $U(1)$ -bundle. The associated bundle is non-trivial since the Chern number is odd, and hence the total space is the dP_1 .

Let us describe the Calabi-Yau metric (1) from the point of view of toric geometry. The metric has an isometry T^3 , locally generated by the Killing vector fields, $\partial/\partial\psi$, $\partial/\partial\phi$ and $\partial/\partial\beta$. The symplectic (Kähler) form ω is given by

$$\omega = \frac{1}{3} \left(d\psi \wedge d(x+z) + d\phi \wedge d((1-x)(1-z)\cos\theta) + d\beta \wedge d(xz) \right). \quad (10)$$

Using the following generators of the T^3 action [10],

$$\frac{\partial}{\partial\phi_1} = \frac{\partial}{\partial\psi} + \frac{\partial}{\partial\phi} - \frac{\partial}{\partial\beta}, \quad (11)$$

$$\frac{\partial}{\partial\phi_2} = \frac{\partial}{\partial\phi} + 3\ell \frac{\partial}{\partial\beta}, \quad (12)$$

$$\frac{\partial}{\partial\phi_3} = -6\ell \frac{\partial}{\partial\beta}, \quad (13)$$

one has Darboux coordinates (ξ^1, ξ^2, ξ^3) on which the symplectic form takes the standard form $\omega = d\xi^i \wedge d\phi_i$:

$$\xi^1 = \frac{1}{3}(1-x)(1-z)(1-\cos\theta), \quad (14)$$

$$\xi^2 = -\frac{\ell}{2}(2xz+1) - \frac{1}{3}(1-x)(1-z)\cos\theta, \quad (15)$$

$$\xi^3 = \ell(2xz+1) \quad (16)$$

with $\ell^{-1} = -5 + 2\sqrt{13}$. For the range of variables: $0 \leq \theta \leq \pi, x_1 \leq x \leq x_2, z \leq z_1$, where x_1, x_2 and z_1 are given by (8), we find the Delzant polytope (see Fig.1)

$$P = \{\xi = (\xi^1, \xi^2, \xi^3) \in \mathbb{R}^3 \mid (\xi, v_a) \geq \lambda_a, a = 1, 2, \dots, 5\}. \quad (17)$$

Here, each v_a is a primitive element of the lattice $\mathbb{Z}^3 \subset \mathbb{R}^3$ and an inward-pointing normal vector to the two dimensional face of P . Explicitly, the set of five vectors $v_a = (1, w_a)$ can be chosen as (see Fig. 2)

$$w_1 = (-1, -2), \quad w_2 = (0, 0), \quad w_3 = (-1, 0), \quad w_4 = (-2, -1), \quad w_5 = (-1, -1). \quad (18)$$

The constants λ_a are given by

$$\lambda_1 = \frac{1}{72}(1 - 5\sqrt{13}), \quad \lambda_3 = \frac{1}{216}(29 + 17\sqrt{13}), \quad \lambda_5 = \frac{1}{54}(31 + 7\sqrt{13}), \quad (19)$$




Figure 1: Delzant polytope P




Figure 2: Toric diagram

and $\lambda_2 = \lambda_4 = 0$. The inner products (ξ, v_a) are evaluated as

$$\begin{aligned}
 (\xi, v_1) &= \frac{2}{9}(1 + \sqrt{13})(x - x_1)(x_1 - z) + \lambda_1, \\
 (\xi, v_2) &= \frac{1}{3}(1 - x)(1 - z)(1 - \cos \theta), \\
 (\xi, v_3) &= \frac{2}{27}(7 + \sqrt{13})(x_2 - x)(x_2 - z) + \lambda_3, \\
 (\xi, v_4) &= \frac{1}{3}(1 - x)(1 - z)(1 + \cos \theta), \\
 (\xi, v_5) &= \frac{2}{27}(-2 + \sqrt{13})(x - z_1)(z_1 - z) + \lambda_5.
 \end{aligned} \tag{20}$$

Thus, we see that the five faces $F_a = \{\xi \in \mathbb{R}^3 \mid (\xi, v_a) = \lambda_a\}$ correspond to degeneration surfaces at $x = x_1$, $\theta = 0$, $x = x_2$, $\theta = \pi$ and $z = z_1$, respectively.

Finally, we note that a D3-brane solution can be constructed from the Calabi-Yau

metric given by g (1) with the special parameters a, b :

$$g^{(10)} = H^{-1/2} g^{(3+1)} + H^{1/2} g, \quad (21)$$

$$F_5 = (1 + *_{10}) \text{vol}_{(3+1)} \wedge dH^{-1}.$$

We find the warp factor H as a harmonic function $\Delta_g H = 0$;

$$H(z) = -\frac{(7 - 2\sqrt{13})L^4}{27} \log \left(\frac{8z^3 - 12z^2 + 137 + 37\sqrt{13}}{(2z + 2 + \sqrt{13})^3} \right) \quad (22)$$

$$-\frac{(10 - 2\sqrt{13})L^4}{27\sqrt{6 + 2\sqrt{13}}} \left(\frac{\pi}{2} - \arctan \left(\frac{-4z + 5 + \sqrt{13}}{3\sqrt{6 + 2\sqrt{13}}} \right) \right), \quad (23)$$

where the constant L is given by

$$L^4 = \frac{4\pi^4 g_s (\alpha')^2 N}{\text{Vol}(Y^{2,1})} = 16(-46 + 13\sqrt{13})\pi g_s (\alpha')^2 N. \quad (24)$$

For large $-z = r^2/2$, the warp factor behaves as

$$H = \frac{L^4}{r^4} + O(1/r^6), \quad (25)$$

while near $z = z_1$

$$H \simeq \frac{2(7 - 2\sqrt{13})L^4}{27} \log(z_1 - z). \quad (26)$$

Acknowledgements

We would like to thank K. Maruyoshi for useful discussions. This work is supported by the 21 COE program ‘‘Constitution of wide-angle mathematical basis focused on knots.’’ The work of Y.Y. is supported by the Grant-in Aid for Scientific Research (No. 17540262 and No. 17540091) from Japan Ministry of Education. The work of T.O. is supported in part by the Grant-in-Aid for Scientific Research (No.18540285) from the Ministry of Education, Science and Culture, Japan.

References

- [1] J. Maldacena, “The large N limit of superconformal field theories and supergravity,” Adv. Theor. Math. Phys. **2** (1998) 231-252; Int. J. Theor. Phys. **38** (1999) 1113-1133, [hep-th/9711200](#);
S.S. Gubser, I.R. Klebanov and A.M. Polyakov, “Gauge theory correlators from non-critical string theory,” Phys. Lett. **B428** (1998) 105-114, [hep-th/9802109](#);
E. Witten, “Anti-de Sitter space and holography,” Adv. Theor. Math. Phys. **2** (1998) 253-291, [hep-th/9802150](#);
O. Aharony, S.S. Gubser, J.M. Maldacena, H. Ooguri and Y. Oz, “Large N field theories, string theory and gravity,” Phys. Rept. **323** (2000) 183-386, [hep-th/9905111](#).
- [2] D.N. Page and C.N. Pope, “Inhomogeneous Einstein Metric On Complex Line Bundles,” Class. Quant. Grav. **4** (1987) 213-225.
- [3] L.A. Pando Zayas and A.A. Tseytlin, “3-Branes on Resolved Conifold,” JHEP **0011** (2000) 028, [hep-th/0010088](#).
- [4] L.A. Pando Zayas and A.A. Tseytlin, “3-Branes on Spaces with $R \times S^2 \times S^3$ Topology,” Phys. Rev. **D63** (2001) 086006, [hep-th/0101043](#).
- [5] S.S. Pal, “A new Ricci flat geometry,” Phys. Lett. **B614** (2005) 201-206, [hep-th/0501012](#).
- [6] K. Sfetsos and D. Zoakos, “Supersymmetric solutions based on $Y^{p,q}$ and $L^{p,q,r}$,” Phys. Lett. **B625** (2005) 135-144, [hep-th/0507169](#).
- [7] S. Benvenuti, M. Mahato, L.A. Pando Zayas and Y. Tachikawa, “The Gauge/Gravity Theory of Blown up Four Cycles,” [hep-th/0512061](#).
- [8] W. Chen, H. Lü and C.N. Pope, “Kerr-de Sitter Black Holes with NUT Charges,” [hep-th/0601002](#);

W. Chen, H. Lü and C.N. Pope, “General Kerr-NUT-AdS Metrics in All Dimensions,” `hep-th/0604125`.

- [9] J.P. Gauntlett, D. Martelli, J. Sparks and D. Waldram, “Sasaki-Einstein metrics on $S^2 \times S^3$,” *Adv. Theor. Math. Phys.* **8** (2004) 711-734, `hep-th/0403002`;
D. Martelli and J. Sparks, “Toric Sasaki-Einstein metrics on $S^2 \times S^3$,” *Phys. Lett.* **B621** (2005) 208-212, `hep-th/0505027`.
- [10] D. Martelli and J. Sparks, “Toric Geometry, Sasaki-Einstein Manifolds and a New Infinite Class of *AdS/CFT* Duals,” *Commun. Math. Phys.* **262** (2006) 51-89, `hep-th/0411238`.